

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

TRAUMA ORGAN PROTECTION WITH ARTESUNATE (TOP-ART): PRECLINICAL EFFICACY OF AN ANTIMALARIAL DRUG WITH EXCELLENT SAFETY PROFILE AND PLANNED PHASE II RCT.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1563297> since 2016-05-29T17:46:24Z

Published version:

DOI:10.1097/01.shk.0000472007.28235.3c

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

HEM-3:

TRAUMA ORGAN PROTECTION WITH ARTESUNATE (TOP-ART): PRECLINICAL EFFICACY OF AN ANTIMALARIAL DRUG WITH EXCELLENT SAFETY PROFILE AND PLANNED PHASE II RCT.

By: Sordi, R; Nandra, K K; Chiazza, F; Patel, N S A; Brohi, K; Collino, M; Thiernemann, C

Shock (Augusta, Ga.)

Volume: 44 Suppl 2

Pages: 1

DOI: 10.1097/01.shk.0000472007.28235.3c

Published: 2015-Oct

Abstract

INTRODUCTION: Hemorrhagic shock (HS) is a common cause of death in severely injured patients and is associated with impairment of organ perfusion, systemic inflammatory response and multiple organ failure. The aim of the present study was to evaluate the effects of artesunate, the drug of choice for the treatment of falciparum malaria, on organ injury and dysfunction associated with HS in the rat.

METHODS: Rats were subjected to HS. The arterial pressure was reduced to 30 mmHg for 90min, followed by resuscitation with shed blood over 5min. Rats were treated with artesunate (2.4 or 4.8mg/kg; i.v.) or vehicle upon resuscitation. Four hours later, organ injury and dysfunction and the signalling events involved in the observed protective effects of artesunate were investigated.

RESULTS: HS resulted in a significant decrease in creatinine clearance, as well as rises in serum creatinine, aspartate aminotransferase, alanine aminotransferase and lactate, indicating the development of renal dysfunction, liver and muscular injury and organ ischemia. HS also caused significant increases in lung myeloperoxidase activity, NF-kappaB activation, iNOS expression and pro-inflammatory cytokines formation. Treatment of HS-rats with artesunate protected animals against the organ injury and dysfunction. Artesunate increased the activation of Akt and eNOS, inhibited the activation of GSK-3beta and NF-kappaB activation, and attenuated the increase in serum TNF-alpha and IL-6 associated with HS.

CONCLUSION: A single-centre placebo-controlled randomized phase II clinical trial will be conducted at Barts Health NHS Trust (TOP-ART) in late 2015 to evaluate the effects of GMP-artesunate in patients with severe hemorrhage following trauma.

Author Information

Address: 1William Harvey Research Institute, Queen Mary University of London, London, UK 2University of Turin, Department of Drug Science and Technology, Turin, Italy 3Blizard Institute, Queen Mary University of London, London, UK.